



**EFFECT OF NURSE LED EDUCATION PROGRAMME ON KNOWLEDGE & EXPRESSED PRACTICE REGARDING RHEUMATOID ARTHRITIS AND ITS SELF-CARE AMONG PATIENTS ATTENDING THE DEPARTMENT OF RHEUMATOLOGY, KGMU**

**Akshita Awasthi<sup>1\*</sup>, Rohitash Kumar<sup>2</sup>, Urmila Dhakad<sup>3</sup>, Puneet Kumar<sup>4</sup> and Prashant Kumar Bajpai<sup>5</sup>**

<sup>1</sup>Nursing Tutor, KGMU, College of Nursing, Lucknow, U.P.

<sup>2</sup>Assistant Professor, KGMU, College of Nursing, Lucknow, U.P.

<sup>3</sup>Head of Department, Department of Rheumatology, KGMU, College of Nursing, Lucknow, U.P.

<sup>4</sup>Associate Professor, Department of Rheumatology, KGMU, Lucknow, U.P.

<sup>5</sup>Assistant Professor, Department of Community Medicine, KGMU, Lucknow, U.P.

Article Received date: 04 April 2025

Article Revised date: 24 April 2025

Article Accepted date: 14 May 2025



\*Corresponding Author: Akshita Awasthi

Nursing Tutor, KGMU, College of Nursing, Lucknow, U.P.

**ABSTRACT**

**Introduction:** Rheumatoid arthritis (RA) is a chronic autoimmune disorder that requires lifelong management, including medication adherence and self-care. Despite medical advancements, many patients lack sufficient knowledge and do not practice effective self-care, leading to disease progression. Nurse-led education programmes can empower patients by improving disease understanding and encouraging proper self-management behaviours. **Methodology:** A quasi-experimental one-group pre-test post-test design was adopted to assess the effect of a nurse-led education programme on knowledge and expressed self-care practices among 92 RA patients attending the Department of Rheumatology, King George's Medical University (KGMU), Lucknow. Data were collected using a socio-demographic questionnaire, a self-structured knowledge questionnaire (18 items), and a self-care practice checklist (15 items). Pre- and post-intervention assessments were conducted one month apart. Data were analyzed using descriptive and inferential statistics, including paired t-tests and chi-square tests. **Results:** The study revealed a significant improvement in participants' knowledge post-intervention. Pre-test results showed 34.8% of patients had poor knowledge, while post-test results indicated that 94.6% had good knowledge ( $p < 0.05$ ). Although an increase in self-care practice scores was noted, the change was not statistically significant. No significant association was found between demographic variables and knowledge or practice scores. **Conclusion:** The nurse-led education programme was effective in improving knowledge among RA patients. However, the intervention had a limited impact on expressed self-care practices, indicating the need for sustained, behaviour-focused interventions in conjunction with education.

**KEYWORDS:** Rheumatoid Arthritis, Nurse-Led Intervention, Patient Education, Self-Care Practices, Knowledge, Chronic Disease Management.

**INTRODUCTION**

Rheumatoid arthritis (RA) is a chronic autoimmune disorder that primarily affects the synovial joints, leading to inflammation, pain, stiffness, and progressive joint damage. The condition significantly impairs the quality of life, affecting mobility and daily activities. Globally, RA affects approximately 0.24 to 1% of the population, with a higher prevalence among women.<sup>[1]</sup> The exact etiology remains unclear, but genetic predisposition, environmental triggers, and immune system dysregulation contribute to its development. Without timely

intervention and proper disease management, RA can lead to severe joint deformities, disability, and systemic complications affecting the cardiovascular, respiratory, and nervous systems. Given its lifelong nature, effective disease management strategies are essential to control disease progression and improve patient outcomes.<sup>[2]</sup>

Patient education is a cornerstone in RA management, as it empowers individuals to understand their condition, adhere to prescribed treatment regimens, and implement self-care strategies. Studies have shown that patients

with higher disease-related knowledge demonstrate better medication adherence, improved physical functioning, and reduced disease flares.<sup>[3]</sup> In many healthcare settings, particularly in resource-limited regions, patient education remains inadequate. Many RA patients rely solely on medications without understanding the importance of lifestyle modifications, joint protection techniques, regular physical activity, and dietary interventions.<sup>[4]</sup> This knowledge gap often results in poor disease control, increased dependency on medical care, and reduced quality of life.

Nurses play a pivotal role in bridging this gap through structured education programmes aimed at improving patients' understanding of RA and its self-care management. Nurse-led education has been recognized as an effective approach in chronic disease management, enhancing patients' knowledge, self-efficacy, and adherence to recommended care plans.<sup>[5]</sup> By delivering disease-specific information, teaching symptom management strategies, and reinforcing the importance of self-care, nurses can significantly improve patient outcomes. This study focuses on evaluating the impact of a structured nurse-led education programme on the knowledge and expressed self-care practices of RA patients, emphasizing the need for continuous education to promote better disease management and overall well-being.

Global prevalence of rheumatoid arthritis (RA) is estimated at approximately 0.46%.<sup>[6]</sup> In India, prevalence estimates vary, with studies indicating rates ranging from 0.28% to 0.70%.<sup>[7]</sup> Notably, a study from Lucknow, Uttar Pradesh, reported a prevalence of 0.19% in urban areas and 0.54% in rural areas.<sup>[8]</sup> These variations underscore the need for region-specific data to accurately assess the burden of RA.

The disease predominantly affects women, with studies indicating that approximately 70% of RA patients are female.<sup>[9]</sup> The onset is most common between the ages of 30 and 60, although it can occur at any age.<sup>[10]</sup> Despite advancements in treatment, a significant proportion of patients lack adequate knowledge about RA management. This knowledge gap often leads to poor adherence to treatment regimens and suboptimal self-care practices, resulting in disease progression and reduced

quality of life.

Nurse-led education programs have been shown to enhance patient knowledge and self-management skills, leading to improved health outcomes. There is a scarcity of structured educational interventions in India, particularly in tertiary healthcare settings. This study aims to assess the effectiveness of a nurse-led education program in improving knowledge and self-care practices among RA patients at King George's Medical University (KGMU), Lucknow, thereby addressing the critical gap between awareness and practical disease management.

## METHODOLOGY

**Research approach:** Quantitative research approach.

**Research design:** A quasi-experimental one-group pre-test post-test design.

**Study Setting and Participants:** The study was conducted in the Department of Rheumatology at King George's Medical University (KGMU), Lucknow, Uttar Pradesh, a well-established tertiary care centre catering to rheumatoid arthritis (RA) patients.

**Sampling:** A non-probability purposive sampling technique was used to select 92 rheumatoid arthritis (RA) patients attending the Department of Rheumatology at KGMU, Lucknow. The sample size was determined based on statistical calculations to ensure adequate representation.

**Data collection:** Data collection involved structured questionnaires and checklists to assess demographics, knowledge, and self-care practices among RA patients. A pre-test was followed by a nurse-led education programme, with a post-test conducted after one month to evaluate improvements.

**Data analysis:** The data analysis involved descriptive statistics (frequency, percentage, mean, standard deviation) to summarize demographics, knowledge, and self-care practices, and inferential statistics (paired t-test, chi-square test) to assess the effectiveness of the nurse-led education programme and associations with socio-demographic variables.

## RESULT

**Table 1: Frequency and Percentage distribution of demographical characteristics in samples. N=92**

| S. No. | Demographic variable | Frequency (f) | Percent (%) |
|--------|----------------------|---------------|-------------|
| 1      | Age in years         |               |             |
|        | a) 18-35 years       | 38            | 41.3        |
|        | b) 36-45 years       | 29            | 31.5        |
|        | c) 46-55 years       | 15            | 16.3        |
|        | d) 56-65 years       | 10            | 10.9        |
| 2      | Gender               |               |             |
|        | a) Male              | 23            | 25.0        |
|        | b) Female            | 69            | 75.0        |
|        | c) Transgender       | 0             | 0           |

|     |   |    |      |
|-----|---|----|------|
| 3   | Religion  |    |      |
|     | a) Hindu  | 80 | 87.0 |
|     | b) Muslim   | 12 | 13.0 |
|     | c) Christian  | 0  | 0    |
| 4   | Educational Status  |    |      |
|     | a) Illiterate   | 11 | 12.0 |
|     | b) Primary education  | 10 | 10.9 |
|     | c) Secondary education  | 24 | 26.1 |
|     | d) Graduate   | 41 | 44.6 |
|     | e) Post Graduate  | 6  | 6.5  |
| 5   | Type of family  |    |      |
|     | a) Nuclear  | 15 | 16.3 |
|     | b) Joint  | 70 | 76.1 |
|     | c) Extended   | 7  | 7.6  |
| 6   | Marital status  |    |      |
|     | a) Married  | 67 | 72.8 |
|     | b) Unmarried  | 21 | 22.8 |
|     | c) Widowed  | 1  | 1.1  |
|     | d) Divorced/Separated   | 3  | 3.3  |
| 7   | Monthly income in rupees  |    |      |
|     | a) Less than Rs. 5000/-   | 4  | 4.3  |
|     | b) Rs. 5001 to Rs. 10000/-  | 14 | 15.2 |
|     | c) More than Rs. 10001/-  | 74 | 80.4 |
| 8   | Occupation  |    |      |
|     | a) Home makers  | 34 | 37.0 |
|     | b) Agriculture  | 11 | 12.0 |
|     | c) Daily wage worker  | 12 | 13.0 |
|     | d) Self employed  | 18 | 19.6 |
|     | e) Private job  | 16 | 17.4 |
|     | f) Government job   | 1  | 1.1  |
| 9   | Numbers of family members   |    |      |
|     | a) Less than 5 members  | 50 | 54.3 |
|     | b) More than 5 members  | 42 | 45.7 |
| 10a | Previous information regarding Rheumatoid arthritis               |    |      |
|     | a) Yes  | 86 | 93.5 |
|     | b) No   | 6  | 6.5  |
| 11  | Source of information regarding Rheumatoid arthritis              |    |      |
|     | a) Health care workers  | 79 | 91.9 |
|     | b) Mass media   | 5  | 5.8  |
|     | c) Family member and peer group                                   | 2  | 2.3  |
| 12  | Previous information regarding self-care of Rheumatoid arthritis  |    |      |
|     | a) Yes  | 16 | 17.4 |
|     | b) No   | 76 | 82.6 |
| 13  | source of information regarding self-care of Rheumatoid arthritis |    |      |
|     | a) Health care workers  | 11 | 68.8 |
|     | b) Mass media   | 5  | 31.3 |

**Table 2: Showing level the Pretest and Posttest knowledge regarding rheumatoid Arthritis and Its self-care among patients attending the department of Rheumatology. N= 92**

| S. No | Level of knowledge | Pretest   |            | Posttest  |            |
|-------|--------------------|-----------|------------|-----------|------------|
|       |                    | Frequency | Percentage | Frequency | Percentage |
| 1.    | Poor knowledge     | 32        | 34.8       | 0         | 0.0        |
| 2.    | Average knowledge  | 39        | 42.4       | 5         | 5.4        |
| 3.    | Good knowledge     | 21        | 22.8       | 87        | 94.6       |

**Table 3: Showed “t” test value for assessment of effectiveness of nurse led education programme on knowledge regarding rheumatoid arthritis and its self-care among patients attending the department of Rheumatology.**

| Test     | Mean  | S.D.  | Mean difference | Calculated “t” value | df | p-value |
|----------|-------|-------|-----------------|----------------------|----|---------|
| Pretest  | 8.00  | 3.190 | 5.50            | 20.66                | 91 | 0.001*  |
| Posttest | 13.50 | 1.788 |                 |                      |    |         |

**Table 4: Showing level the pre-test and post-test expressed practice regarding rheumatoid arthritis and its self-care among patients attending the department of Rheumatology. N= 92**

| S. No | Level of expressed practice | Pretest   |            | Posttest  |            |
|-------|-----------------------------|-----------|------------|-----------|------------|
|       |                             | Frequency | Percentage | Frequency | Percentage |
| 1.    | Poor self-care              | 32        | 34.8       | 6         | 6.5        |
| 2.    | Average self-care           | 49        | 53.3       | 81        | 88.0       |
| 3.    | Good selfcare               | 11        | 12.0       | 5         | 5.4        |

**Table 5: Chi Square value showing association between pretest knowledge regarding rheumatoid arthritis and its self-care among patients attending the department of Rheumatology and their selected demographic variables N=92**

| S. N. | Name of variables  | Chi square | df | “p” value |
|-------|--|------------|----|-----------|
| 1     | Age in years   | 11.093     | 6  | 0.086     |
| 2     | Gender   | 0.396      | 2  | 0.82      |
| 3     | Religion   | 0.419      | 2  | 0.811     |
| 4     | Educational Status   | 12.892     | 8  | 0.116     |
| 5     | Type of family   | 2.267      | 4  | 0.687     |
| 6     | Marital status   | 4.742      | 6  | 0.577     |
| 7     | Monthly income in rupees   | 3.328      | 4  | 0.505     |
| 8     | Occupation   | 10.06      | 10 | 0.435     |
| 9     | Numbers of family members  | 3.526      | 2  | 0.172     |
| 10    | Previous information regarding Rheumatoid arthritis              | 20.77      | 2  | 0.001*    |
| 12    | Previous information regarding self-care of Rheumatoid arthritis | 5.992      | 2  | 0.049*    |

\*= Significant

NS = Non-significant

## DISCUSSION

The present study was conducted to assess the effectiveness of a nurse-led education programme on knowledge and expressed self-care practices among rheumatoid arthritis (RA) patients. The results clearly indicate that the educational intervention had a significant positive impact on the patients' knowledge levels, with 94.6% of participants achieving good knowledge in the post-test compared to only 22.8% in the pre-test. While knowledge improved significantly, the correlation between knowledge and expressed self-care practice was not statistically significant, suggesting that increased knowledge does not always translate into behavioural change. This highlights a critical gap in health education where knowledge alone may not ensure proper self-management.

In our study, the implementation of a nurse-led education program resulted in a significant increase in knowledge among RA patients, with the proportion of participants displaying good knowledge rising from 22.8% pre-intervention to 94.6% post-intervention. This outcome mirrors findings from Gurjar (2023), who demonstrated that a structured, nurse-led educational program

significantly improved both knowledge and self-care behaviour among arthritis patients after a 3-month intervention.<sup>[11]</sup> Similarly, Selim et al. (2019) observed significant improvements in knowledge scores among elderly RA patients following nursing interventions, confirming that targeted education is vital to empowering patients.<sup>[12]</sup>

Regarding self-care practices, our results showed that poor self-care dropped from 34.8% to just 6.5% post-intervention, while average self-care increased dramatically. This aligns with the findings from Ndosi (2016), who highlighted that combining nurse-led care with patient self-assessment enhanced disease control and management of RA-related comorbidities.<sup>[13]</sup> Yang et al. (2025) further supported this by reporting that nurse-led care significantly improved disease activity and self-efficacy compared to usual care.<sup>[14]</sup> These findings support the idea that knowledge gained through education directly translates into improved health behaviours, particularly in chronic disease management.

Our study also found significant associations between patients' prior knowledge and their pre-test scores, with

*p*-values of 0.001 and 0.049 respectively. This indicates that patients with baseline knowledge were more likely to benefit from the intervention. This trend is consistent with a systematic review by Lois et al. (2021), which emphasized that nurse-led care enhances treatment adherence, depression outcomes, and overall patient satisfaction through sustained engagement and support.<sup>[15]</sup> Garner et al. (2017) confirmed that nurse-led care improved patient satisfaction and was perceived as effective and safe across multiple quality-of-care domains.<sup>[16]</sup>

One of the strengths of the current study is the implementation of a structured nurse-led education programme within a tertiary care setting, using validated tools for assessment. This intervention was found to be both feasible and effective in significantly improving knowledge levels. However, the study also had limitations. It lacked a control group, which limits the ability to establish causality. The sample was selected through purposive sampling, which may affect the generalizability of the findings. The follow-up period was short (only one month), which may not have been sufficient to capture changes in long-term practice or behavior. An unexpected finding was the minimal improvement in self-care practices despite the significant increase in knowledge—indicating a need to integrate behavioral training or motivational strategies in future interventions.

The primary hypothesis of the study—that there would be a significant difference between pre- and post-test knowledge and practice scores among RA patients following a nurse-led education programme—was partially supported. The knowledge scores showed a statistically significant improvement, while the change in expressed practice scores was less pronounced, highlighting a gap between learning and application. The study's purpose of evaluating the impact of a nurse-led intervention was achieved, demonstrating that structured educational programmes can effectively raise awareness and understanding of RA among patients.

This study highlights the significance of nurse-led interventions in chronic disease management and suggests the importance of continuous reinforcement to achieve behavioural change. Future research should focus on incorporating motivational interviewing, family involvement, and psychosocial support to enhance self-care adherence. Additionally, longer follow-up periods and multi-centre randomized controlled trials could provide stronger evidence and help develop more sustainable, culturally tailored education models for RA and other chronic conditions.

## REFERENCES

1. Venetsanopoulou AI, Alamanos Y, Voulgari PV, Drosos AA. Epidemiology and Risk Factors for Rheumatoid Arthritis Development. *Mediterr J Rheumatol*, 2023; 30, 34(4): 404–13.
2. Wu D, Luo Y, Li T, Zhao X, Lv T, Fang G, et al. Systemic complications of rheumatoid arthritis: Focus on pathogenesis and treatment. *Front Immunol*, 2022; 22, 13: 1051082.
3. Marengo MF, Suarez-Almazor ME. Improving treatment adherence in patients with rheumatoid arthritis: what are the options? *Int J Clin Rheumatol*, 2015; 1, 10(5): 345–56.
4. Schäfer C, Keyßer G. Lifestyle Factors and Their Influence on Rheumatoid Arthritis: A Narrative Review. *J Clin Med*, 2022; 2, 11(23): 7179.
5. Berardinelli D, Conti A, Hasnaoui A, Casabona E, Martin B, Campagna S, et al. Nurse-Led Interventions for Improving Medication Adherence in Chronic Diseases: A Systematic Review. *Healthcare*, 2024; 22, 12(23): 2337.
6. Almutairi K, Nossent J, Preen D, Keen H, Inderjeeth C. The global prevalence of rheumatoid arthritis: a meta-analysis based on a systematic review. *Rheumatol Int*, 2021; 41(5): 863–77.
7. Shi W, Liang X, Zhang H, Li H. Burden of rheumatoid arthritis in India from 1990 to 2021: insights from the Global Burden of Disease Database. *Front Med [Internet]*, 2025; 14 [2025, 17]: 12. Available from: <https://www.frontiersin.org/journals/medicine/articles/10.3389/fmed.2025.1526218/full>
8. Misra DP, Sharma A, Dharmanand BG, Chandrashekara S. The Epidemiology of Rheumatic Diseases in India. *Indian J Rheumatol*, 2024; 1, 19(1): 54–61.
9. Rheumatoid arthritis [Internet]. [2025; 17]. Available from: <https://www.who.int/news-room/fact-sheets/detail/rheumatoid-arthritis>
10. Pavlov-Dolijanovic S, Bogojevic M, Nozica-Radulovic T, Radunovic G, Mujovic N. Elderly-Onset Rheumatoid Arthritis: Characteristics and Treatment Options. *Medicina (Mex)*, 2023; 23, 59(10): 1878.
11. Gurjar NR. Effect of educational program on knowledge and self-care behavior among arthritis patients: Pre-experimental research design. *Int J Orthop Trauma Nurs*, 2024; 1, 52: 101038.
12. Selim T, Elashri N, Hasanein AA. Effect of Nursing Interventions on Self Care Behaviors and Knowledge for Rheumatoid Arthritic Elderly Patients. In, 2019; [2025, 21]. Available from: [https://www.semanticscholar.org/paper/Effect-of-Nursing-Interventions-on-Self-Care-and-Selim-Elashri/49be11d999726d57a3d5fe435dc6432d518498a6?utm\\_source=consensus](https://www.semanticscholar.org/paper/Effect-of-Nursing-Interventions-on-Self-Care-and-Selim-Elashri/49be11d999726d57a3d5fe435dc6432d518498a6?utm_source=consensus)
13. Ndosi M. Combining nurse-led care with patient self-assessment could enhance tight control of rheumatoid arthritis and management of comorbidities. *Evid Based Nurs*, 2016; 1, 19(3): 85–6.
14. Yang L, Xiang P, Pi G, Wen T, Liu L, Liu D. Effectiveness of nurse-led care in patients with rheumatoid arthritis: a systematic review and meta-analysis. *BMJ Open Qual [Internet]*, 2025; 4 [2025,

- 21]: 14(1). Available from:  
<https://bmjopenquality.bmj.com/content/14/1/e003037>
15. Lois P, Pedraza LL, Candelas G, Lajas C, Rodero M, López TM, et al. AB0906-HPR BENEFIT OF A NURSE-LED PROGRAM OF CARE FOR MANAGEMENT OF PATIENTS WITH ESTABLISHED RHEUMATOID ARTHRITIS: SYSTEMATIC LITERATURE REVIEW. *Ann Rheum Dis*, 2021; 1, 80: 1475.
  16. Garner S, Lopatina E, Rankin JA, Marshall DA. Nurse-led Care for Patients with Rheumatoid Arthritis: A Systematic Review of the Effect on Quality of Care. *J Rheumatol*, 2017; 1, 44(6): 757–65.